

# Genetic diversity of *Trichoderma atroviride* strains collected in Poland and identification of loci useful in detection of within-species diversity

Dominik Skoneczny · Michał Oskiera ·  
Magdalena Szczech · Grzegorz Bartoszewski

Received: 19 August 2014 / Accepted: 5 March 2015 / Published online: 20 March 2015  
© The Author(s) 2015. This article is published with open access at Springerlink.com

**Abstract** Molecular markers that enable monitoring of fungi in their natural environment or assist in the identification of specific strains would facilitate *Trichoderma* utilization, particularly as an agricultural biocontrol agent (BCA). In this study, sequence analysis of internal transcribed spacer regions 1 and 2 (ITS1 and ITS2) of the ribosomal RNA (rRNA) gene cluster, a fragment of the translation elongation factor 1- $\alpha$  (*tef1*) gene, and random amplified polymorphic DNA (RAPD) markers were applied to determine the genetic diversity of *Trichoderma atroviride* strains collected in Poland, and also in order to identify loci and PCR-based molecular markers useful in genetic variation assessment of that fungus. Although *tef1* and RAPD analysis showed limited genetic diversity among *T. atroviride* strains collected in Poland, it was possible to distinguish major groups that clustered most of the analyzed strains. Polymorphic RAPD amplicons were cloned and sequenced, yielding sequences representing 13 *T. atroviride* loci. Based on these sequences, a set of PCR-based markers specific to *T. atroviride* was developed and examined. Three cleaved amplified polymorphic sequence (CAPS) markers could assist in distinguishing *T. atroviride*

strains. The genomic regions identified may be useful for further exploration and development of more precise markers suitable for *T. atroviride* identification and monitoring, especially in environmental samples.

## Introduction

*Trichoderma* is a fungal genus that includes more than 200 species that occur all over the world in different geographical regions and climatic zones (Atanasova et al. 2013; Hoyos-Carvajal et al. 2009; Kredics et al. 2014). They are remarkable for their rapid growth under various environmental conditions, as well as their ability to use different substrates as carbon sources (Kubicek et al. 2003; Atanasova and Druzhinina 2010); thus, they are recognized as efficient decomposers of woody and herbal materials and other organic matter (Atanasova et al. 2013; Schuster and Schmoll 2010). *Trichoderma* interacts with other microorganisms and plants in diverse ways (Harman et al. 2004; Lorito et al. 2010; Druzhinina et al. 2011; Woo et al. 2006). *Trichoderma* strains, including *Trichoderma atroviride* strains, that are promoting efficient plant growth and stress resistance are used as biological control agents (BCAs) in sustainable farming systems (Chet and Inbar 1994; Benítez et al. 2004; Stewart and Hill 2014).

Various molecular techniques have been used in *Trichoderma* research, in order to investigate genetic diversity within the genus. These include restriction fragment length polymorphism (RFLP) analysis (Dodd et al. 2004a), random amplified polymorphic DNA (RAPD) analysis (Hermosa et al. 2001), amplified fragment length polymorphism (AFLP) analysis (Buhariwalla et al. 2005), sequence-characterized amplified region (SCAR) analysis (Hermosa

Dominik Skoneczny and Michał Oskiera contributed equally to this work.

D. Skoneczny · G. Bartoszewski (✉)  
Department of Plant Genetics Breeding and Biotechnology, Warsaw  
University of Life Sciences (SGGW), Nowoursynowska 159,  
02-776 Warszawa, Poland  
e-mail: grzegorz\_bartoszewski@sggw.pl

M. Oskiera · M. Szczech  
Microbiology Laboratory, Research Institute of Horticulture,  
Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland

### Present Address:

D. Skoneczny  
School of Agricultural and Wine Sciences, Charles Sturt University,  
Locked Bag 588, Wagga Wagga, NSW 2678, Australia

et al. 2001; Dodd et al. 2004b; Cordier et al. 2007), microsatellite markers (Naef et al. 2006), and sequence analysis (Kindermann et al. 1998; Dodd et al. 2003; Druzhinina et al. 2005; Błaszczyk et al. 2011; Atanasova et al. 2013). One of the simplest and cheapest techniques is RAPD, which is characterized by low repeatability and transferability between different laboratories. Nevertheless, it delivers solid data at minimal expense when used strictly in a single laboratory. Moreover, RAPD markers can be easily transformed into PCR-based SCAR markers, as demonstrated in several studies for *T. atroviride* (Hermosa et al. 2001; Cordier et al. 2007; Feng et al. 2011).

Application of molecular techniques in *Trichoderma* taxonomy solved inaccuracies related to phenotypical *T. atroviride* species identification, as well as demonstrated interspecies genetic variability. In order to reevaluate biocontrol strains reported as *Trichoderma harzianum* and *Trichoderma viride*, Hermosa et al. (2000) utilized sequences of internal transcribed spacer regions 1 and 2 (ITS1 and ITS2) of the ribosomal RNA (rRNA) gene cluster and analyzed the hybridization patterns of mitochondrial DNA. Authors identified four groups of *Trichoderma* including *T. atroviride*. Identification of mycoparasitic strains performed with assistance of ITS1, ITS2, mtSSUrDNA, and partial *chi18-5* gene sequences revealed that the strains belonged to *T. harzianum*, *T. atroviride*, and *Trichoderma asperellum* (Kullnig et al. 2001). Based on the sequences of ITS1, ITS2, and translation elongation factor 1-alpha (*tef1*) gene fragments, four major clades (A, B, C, and D) of *T. atroviride* were distinguished, and the fifth clade E was suggested (Dodd et al. 2003; Jaklitsch et al. 2006; Samuels et al. 2006; Mulaw et al. 2010; Gal-Hemed et al. 2011).

Application of *Trichoderma* BCA strains raised questions about fungal survival and spread in the environment (Dodd et al. 2004a, b). The molecular strategies were applied to develop markers useful for *Trichoderma* monitoring at different levels: genus-specific (Hagn et al. 2007; Devi et al. 2011; Chakraborty et al. 2011; Friedl and Druzhinina 2012), species-specific (Chen et al. 1999; Kredics et al. 2009; Miyazaki et al. 2009; Friedl and Druzhinina 2012), and strain-specific (Hermosa et al. 2001; Rubio et al. 2005; Cordier et al. 2007; Savazzini et al. 2008). Strain-specific molecular markers proposed for *Trichoderma* BCA strain monitoring included RAPD, RFLP (Zimand et al. 1994; Bowen et al. 1996; Abbasi et al. 1999), and SCAR markers (Hermosa et al. 2001; Dodd et al. 2004b; Rubio et al. 2005; Cordier et al. 2007; Savazzini et al. 2008; Longa et al. 2009; Feng et al. 2011; Naeimi et al. 2011). Due to the complex taxonomy of *Trichoderma*, development of strain-specific markers is a long and laborious process (Druzhinina and Kubicek 2005). The most beneficial is development of species-specific markers, which can also be used to simplify and assist in taxonomic identification, based on sequence

differences at the interspecies level, useful for elaboration of such markers (Chen et al. 1999; Friedl and Druzhinina 2012; Kredics et al. 2009; Miyazaki et al. 2009).

The aim of this study was to utilize sequences of ITS1 and ITS2 region, a fragment of the *tef1* gene, and RAPD markers to analyze the genetic diversity of *T. atroviride* strains collected in Poland, develop PCR-based markers, and identify genomic regions to enable genetic diversity studies and further development of reliable methods of *T. atroviride* identification. In this study, RAPD markers were converted into PCR-based markers suitable for detection of *T. atroviride*. Additionally, application of CAPS markers developed in this study is expected to improve the accuracy of *T. atroviride* classification.

## Materials and methods

### Fungal material and strain identification

All *Trichoderma* strains used in this study were obtained from a stock culture collection maintained by the Microbiology Laboratory at the Research Institute of Horticulture in Skierniewice, Poland. Stock cultures were deep-frozen in glycerol and stored at  $-80^{\circ}\text{C}$ . Forty strains of *T. atroviride* were used in this study; these included 38 strains of *T. atroviride* collected at different locations in Poland, and the two reference strains CBS 693.94 (Dodd et al. 2003) and IMI206040 (Kubicek et al. 2011) (Table 1). Strain identification was based on the internal transcribed spacer regions 1 and 2 (ITS1 and ITS2) of the rRNA gene cluster sequences and *TrichOKey* v. 2.0 (Druzhinina et al. 2005) and sequences of the fragment of the gene encoding translation elongation factor 1-alpha (*tef1*) and *TrichoBLAST* v. 1.0 analysis (Kopchinskiy et al. 2005). All sequences were obtained by PCR and direct amplicon sequencing from both directions. ITS1 and ITS2 regions were PCR-amplified with ITS6 and ITS4 primers (Cooke and Duncan 1997; White et al. 1990). For the *tef1* sequencing, primers EF1-728 F and TEFLLErev were used (Carbone and Kohn 1999; Jaklitsch et al. 2005). ITS and *tef1* sequences of the strains are available in the NCBI GenBank (Accession numbers are provided in Table 1). The following strains were used for the PCR-based marker species specificity evaluation: *Trichoderma aggressivum* f. *aggressivum* CBS 100528, *T. aggressivum* f. *europaeum* CBS 100526, *T. asperellum* TRS705, *Trichoderma citrinoviride* TRS119, *Trichoderma cerinum* 38.24.06.2, *Trichoderma crassum* TRS113, *Trichoderma gamsii* TRS123, *Trichoderma hamatum* TRS127, *T. harzianum* sensu lato II subclade CBS 466.94, *T. harzianum* s. l. III subclade TRS60, *T. harzianum* s. l. X subclade CBS 115901, *T. harzianum* s. l. Lixii subclade TRS66, *T. harzianum* sensu stricto TRS71, *Trichoderma longibrachiatum* TRS708,

**Table 1** Source and origin of *T. atroviride* strains used in this study

Strain	GenBank accession number of sequences		Source	Location	Origin
	ITS	<i>tefl</i>			
TRS31	KJ786740	KJ786821	Soil	Balcerów	Central Poland
TRS15	KJ786724	KJ786805	Lumberyard	Głuchów	
TRS36	KJ786729	KJ786810	Soil	Kątne	
TRS28	KJ786721	KJ786802	Production hall	Kolonia Bolimowska	
TRS5	KJ786741	KJ786822	Compost	Łódź	
TRS2	KJ786749	KJ786830	Compost	Łódź	
TRS1	KJ786739	KJ786820	Mushroom growing hall	Maków	
TRS3	KJ786743	KJ786824	Mushroom growing hall	Maków	
TRS9	KJ786738	KJ786819	Mushroom growing hall	Maków	
TRS18	KJ786757	KJ786839	Compost	Maków	
TRS32	KJ786744	KJ786825	Production hall	Maków	Eastern Poland
TRS45	KJ786718	KJ786799	Compost	Maków	
TRS20 <sup>a</sup>	KJ786730	KJ786811	Production hall	Sierakowice Lewe	
TRS30	KJ786720	KJ786801	Soil	Skiemiewice	
TRS14 <sup>a</sup>	KJ786725	KJ786806	Compost	Skiemiewice	
35.24.06.4 <sup>a</sup>	KJ786742	KJ786823	Field soil	Skiemiewice	
TRS68	KJ786745	KJ786826	Mushroom growing hall	Skiemiewice	
TRS17	KJ786734	KJ786815	Mushroom growing hall	Skiemiewice	
TRS21	KJ786727	KJ786808	Soil	Topołowo	
TRS22	KJ786726	KJ786807	Soil	Wiskitki	
TRS24	KJ786722	KJ786803	Production hall	Iganie Nowe	Northern Poland
TRS25 <sup>a</sup>	KJ786731	KJ786812	Production hall	Iganie Nowe	
TRS38	KJ786733	KJ786814	Soil	Myślubórz	Southern Poland
TRS39	KJ786728	KJ786809	Soil	Myślubórz	
TRS42	KJ786752	KJ786833	Soil	Radzanów	Western Poland
TRS7 <sup>a,b</sup>	KJ786747	KJ786828	Production hall	Bieruń	
TRS40 <sup>a</sup>	KJ786755	KJ786836	Compost	Rzeczyce	Poland
TRS26	KJ786751	KJ786832	Soil	Lipniak	
TRS6	KJ786750	KJ786831	Compost	Sława	Poland
TRS10	KJ786754	KJ786835	Not known	Turek	
TRS43 <sup>a</sup>	KJ786748	KJ786829	Forest soil	Wielkopolski Park Narodowy	Poland
TRS11	KJ786735	KJ786816	Not known	Wrocław	
TRS12 <sup>a</sup>	KJ786737	KJ786818	Not known	Wrocław	Poland
TRS13	KJ786736	KJ786817	Not known	Wrocław	
TRS16	KJ786753	KJ786834	Not known	Not known	Poland
TRS19	KJ786732	KJ786813	Soil	Not known	
TRS34	KJ786719	KJ786800	Not known	Not known	Poland
TRS44	KJ786756	KJ786837	Not known	Not known	
CBS 693.94 (IMI 359825)	AF359263	KJ786838	Mushroom compost	<i>T. atroviride</i> reference strain <sup>c</sup>	Northern Ireland
IMI206040	AF278795.1	ABDG00000000.2	Biocontrol strain	<i>T. atroviride</i> reference strain <sup>d</sup>	USA

<sup>a</sup> Strains used in RAPD primer screening part of study<sup>b</sup> Not included in main genetic diversity study<sup>c</sup> Dodd et al. (2003)<sup>d</sup> Kubicek et al. (2011)

*Trichoderma pleuroticola* TRS120, *Trichoderma spirale* TRS111, *Trichoderma tomentosum* TRS82, *Trichoderma virens* Gv29-8, *Trichoderma viride* TRS575, *Trichoderma*

*viridescens* TRS35, and *Trichoderma velutinum* 29.24.06.1. For all above mentioned strains, species/subclade affiliation was confirmed by ITS and *tefl* sequencing as described above

for *T. atroviride*. Mycelia taken from stock cultures were placed on potato dextrose agar (PDA) plates (Sigma-Aldrich, St. Louis, MO, USA) and cultured in the dark at 25 °C for 4–7 days. Mycelia grown on PDA were then used to inoculate 250-mL flasks containing 50 mL potato dextrose broth (Sigma-Aldrich), and cultures were incubated in the dark on a rotary shaker at 200 rpm for 7 days at room temperature ( $21 \pm 2$  °C).

#### DNA extraction

Total fungal DNA was isolated using the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany), according to the manufacturer's instructions with slight modifications. Fungal cells of tested *Trichoderma* strains were harvested by centrifugation of liquid cultures and dried on filter paper at room temperature ( $21 \pm 2$  °C), prior to grinding in liquid nitrogen with quartz sand (Sigma-Aldrich). Proteinase K and RNase A were added to the lysis buffer. The purity and quantity of isolated DNA were evaluated by agarose gel electrophoresis followed by ethidium bromide staining, and by determination of  $A_{260}/A_{280}$  using a NanoDrop 8000 spectrophotometer (Thermo Scientific, Waltham, MA, USA). Finally, the DNA concentration was adjusted to 1 ng/ $\mu$ L by diluting the samples with 10 mmol/L Tris-HCl, pH 8.0.

#### RAPD analysis

A set of 520 decamer primers (primer kits OPA to OPZ from Operon Technology, Alameda, CA, USA) was used for initial RAPD screening of eight strains of *T. atroviride*: 35.24.06.4, TRS7, TRS12, TRS14, TRS20, TRS25, TRS40, and TRS43. Then, 68 primers that had given the best results were further tested on 39 strains to assess the genetic diversity of *T. atroviride* strains collected in Poland. RAPD-PCR was performed as described by Olczak-Woltman et al. (2009). Reactions contained 1  $\times$  Green PCR Buffer containing 2 mmol/L  $MgCl_2$ , 0.2 mmol/L each dNTP, 1  $\mu$ g bovine serum albumin, 0.5 U *Taq* polymerase (Fermentas, Vilnius, Lithuania), 1  $\mu$ mol/L arbitrary primer, and 2 ng DNA template. PCR was carried out under conditions of initial incubation at 95 °C for 1 min, followed by 10 cycles of 95 °C for 5 s, 37 °C for 30 s, and 72 °C for 30 s; 35 cycles of 95 °C for 5 s, 37 °C for 30 s, and 72 °C for 60 s; and a final extension at 72 °C for 7 min. RAPD-PCR was carried out in a PTC-200 Thermal Cycler (Bio-Rad, Hercules, CA, USA). Amplified DNA fragments were separated by electrophoresis with 2 % agarose–0.5  $\times$  Tris-borate-EDTA (TBE) gels containing ethidium bromide and visualized under UV light. Gel images were edited in CorelDRAW version 12 (Corel Corporation, Ottawa, Ontario, Canada). The binary matrices were constructed based on evaluation of clear RAPD amplicons. Amplicons were scored for presence (1) or absence (0) for

each strain. Cluster analysis of the binary data was performed using the NTSYS-pc v.2.1 program (Exeter Software, Setauket, NY, USA). Similarity matrices were generated using Jaccard's coefficients, and an unweighted pair-group method using arithmetic averages (UPGMA) was chosen to generate the dendrogram from RAPD similarity matrices. To determine how faithfully the dendrogram preserves pairwise distances between original unmolded data, the cophenetic correlation coefficient parameter ( $r$ ) was calculated.

#### Cloning and sequencing

Polymorphic RAPD amplicons were excised from the gel, and DNA was purified using the QIAquick Gel Extraction Kit (Qiagen) according to the manufacturer's protocol. Next, purified amplicons were reamplified under the conditions described for RAPD-PCR, resolved using a 1 % agarose–0.5  $\times$  TBE gel, excised from the gel, and purified. Efficiently reamplified RAPD amplicons were cloned using a TOPO TA Cloning Kit (Invitrogen, Carlsbad, CA, USA). Recombinant plasmids were isolated, and two independent plasmids were used for sequencing of each amplicon (with the exception of Q01-420, for which only one clone was available) using universal M13 forward and reverse primers. The Sequencer 5.1 program (Gene Codes Corporation, Ann Arbor, MI, USA) was used to assemble single reads to obtain consensus sequences in FASTA format. The consensus sequences were deposited at the National Center for Biotechnology Information (NCBI), and GenBank accession numbers are summarized in Table 2 (RAPD-PCR amplicons) and Table 3 (CAPS amplicons).

#### Bioinformatics analysis

For phylogenetic reconstruction, sequences of the fragment of *tef1* gene were assembled with ClustalX (Thompson et al. 1997) and manually edited in CLC Genomic Workbench 7.5 (CLCBio, Aarhus, Denmark), and final block alignment was prepared with a less stringent selection option using GBlocksServer ([http://molevol.cmima.csic.es/castresana/Gblocks\\_server.html](http://molevol.cmima.csic.es/castresana/Gblocks_server.html)). The unconstrained GTR+I+G substitution model, estimated nucleotide frequencies, and substitution values were selected according to Samuels et al. (2006). Metropolis-coupled Markov chain Monte Carlo (MCMCMC) analysis was performed with two runs for one million generations with four chains, with the heating coefficient  $\lambda=0.1$  with MrBayes 3.2.2  $\times$  64 (Ronquist et al. 2012).

RAPD amplicon sequences were compared using basic local alignment search tool (BLAST) (Altschul et al. 1997) with publicly available sequences deposited at both NCBI and the Genome Portal of the Joint Genome Institute (JGI, Grigoriev et al. 2012). At NCBI, blastn and blastx were used



**Table 2** GenBank accession numbers and results of bioinformatics analysis of RAPD-PCR amplicon sequences

RAPD-PCR amplicon	NCBI GenBank Acc. No.	<i>T. atroviride</i> IMI 206040 v.2 unmasked genome BLAST hits (not filtered for low complexity)		<i>T. atroviride</i> IMI 206040 gene catalog (transcripts)		Protein NCBI BLAST hits (non-redundant database)		Pfam domains (database v. 27)
		scaffold	e-value	Similarity	e-value	Similarity	e-value	
A10-933	KF364331	contig_23	0.0	Triat2:317841	9.14E-8	NS	NA	Not found
B07-500	KF364332	contig_29	0.0	Triat2:302951	1.17E-132	Hypothetical protein	7e-42	Not found
E03-391	KF364333	contig_28	0.0	Not found	NA	NS	NA	Not found
E04-458	KF364334	contig_27	0.0	Not found	NA	NS	NA	Not found
H08-917	KF364335	contig_16	0.0	Triat2:254812	0.0	Tetratricopeptide (TPR) protein	8e-116	Tetratricopeptide repeat, PF13414 and PF07719
M06-768	KF364336	contig_22	0.0	Triat2:299234	9.13E-176	NS	NA	Not found
Q01-420	KF364337	contig_26	0.0	Not found	NA	NS	NA	Not found
Q01-885	KF364338	contig_24	0.0	Not found	NA	NS	NA	Not found
R08-1283	KF364339	contig_29	5.06e-53	Triat2:231658	5.05E-64	Putative ankyrin repeat-containing protein	1e-175	Ankyrin repeats (3 copies) PF12796
T07-429	KF364340	contig_27	0.0	Triat2:321796	6.01E-157	Putative ankyrin repeat-containing protein	7e-429	Not found
X16-1030	KF364341	contig_14	0.0	Triat2:172973	0.0	Hypothetical protein	3e-142	Magnesium transporter NIPA PF05653
X18-634	KF364342	contig_23	0.0	Triat2:299500	0.0	Hypothetical protein	3e-35	Not found
Z04-873	KF364343	contig_20	0.0	Triat2:291123	0.0	Hypothetical protein	6e-54	Not found

NS no significant similarity, NA not applicable

to explore the non-redundant protein sequence, nucleotide collection, and high-throughput genomic sequence databases. At JGI, sequences were aligned using blastn to the *T. atroviride* IMI206040 genome version 2 unmasked assemblies. Proteins containing amino acid sequences similar to those encoded by RAPD amplicons were collected and further analyzed using CLC Genomics Workbench 7.5 (CLCBio), and Pfam database search (Punta et al. 2012). RAPD sequences were scanned for sequence repeats using

RepeatMasker (<http://www.repeatmasker.org/>) and mfold (Zuker 2003).

Cleaved amplified polymorphic sequence marker development

The consensus RAPD sequences were mapped on the *T. atroviride* IMI 206040 v. 2 genome sequence deposited at JGI (<http://genome.jgi-psf.org/Triat2/Triat2.home.html>).

**Table 3** Description of CAPS markers and reaction conditions

CAPS amplicon	Primer names and sequences <sup>a</sup> (5'-3')	Amplicon length (bp) <sup>b</sup>	Optimized PCR conditions (annealing temperature/number of cycles)	Restriction enzyme	GenBank Accession number/strain
Q01-4	Q01_4F-GCACACCAACTGCTGGAGCTT Q01_4R-CACGCTGACAATGACCGACAC	1017	66 °C/27 cycles	<i>BsiI</i>	KF576213/TRS15 KF576210/TRS22 KF576211/TRS40
X18-35	X18_3F-AGGCACAGTCCCCTGTTTAGT X18_5R-TGACGATCCTGGTAAGGTTTG	358	65 °C/35 cycles	<i>TaqI</i>	KF576212/TRS36 KF576213/TRS42
Z04-2	Z04_2F-TTACCCAGTGCGGAATCCAAA Z04_2R-TATACGGCGCCTTCCACATTG	1450	66 °C/27 cycles	<i>DraI</i>	KF576214/CBS 693.94 KF576215/TRS45

CAPS cleaved amplified polymorphic sequence

<sup>a</sup> Sequences of PCR amplicons of representative strains were deposited at NCBI

<sup>b</sup> Amplicon length is based on genome sequence of *T. atroviride* reference strain IMI20604

Several sets of primers were designed to amplify RAPD regions or extended RAPD regions based on genome sequence. A standard PCR was performed using *Taq* polymerase (Fermentas) according to the manufacturer's instructions. PCR was carried out under conditions of initial incubation at 94 °C for 3 min, followed by 35 cycles of 94 °C for 30 s, optimized annealing temperature or gradient temperature for 30 s, and 72 °C for 2 min, followed by a final extension at 72 °C for 10 min. The amplified products for selected *T. atroviride* strains were analyzed on agarose gels, purified, and sequenced. Derived sequences were compared and searched for polymorphic restriction sites using Sequencher 5.1 (Gene Codes). PCR amplification was performed using selected primer pairs. PCR products were digested with corresponding restriction enzyme (*Bsi*I, *Dra*I, and *Taq*I; all from Fermentas) according to the manufacturer's instructions, and digestion products were separated on 1.5 % agarose–0.5× TBE gel. The respective primer pairs with optimized PCR conditions and corresponding restriction enzymes are shown in Table 3. To confirm specificity of the CAPS PCR to *T. atroviride* species, multiplex PCR was performed with DNA of 19 *Trichoderma* species mentioned above, CAPS primers, and additional primers 5.8S-R and LR6 (Vilgalys and Hester 1990). The pair of primers 5.8S-R and LR6 amplified a 1.45-kb fragment of the ITS region and served as a positive control for the PCR.

## Results and discussion


Molecular identification of strains based on sequences of ITS1, ITS2, and the fragment of *tef1* gene confirmed that all of the strains used in this study belong to *T. atroviride*. Based on *tef1* sequences, all strains were classified to clade A described by Dodd et al. (2003), except for TRS18 that did not fit any clade. *TrichOKEY* 2 analysis revealed one new ITS1 sequence variant that possesses a single nucleotide change in comparison to the IMI206040 sequence. Six strains, TRS15, TRS24, TRS28, TRS30, TRS34, and TRS45, were characterized by the new ITS1 variant. Seven variants of *tef1* sequence which differed at least by a single nucleotide were found within the studied *T. atroviride* strains. Bayesian phylogeny analysis of *tef1* sequences allowed strain clustering (Fig. 1). TRS18 strain was found to be the most distant one. Four groups of strains were distinguished, with the two biggest groups differing within by a single nucleotide change only. Strains TRS43 and CBS 693.94 did not fall into any groups (Fig. 1).

In addition, RAPD markers were used for further evaluation of genetic diversity of *T. atroviride* strains. A total of 520 primers were tested on eight *T. atroviride* strains, including 35.24.06.4, TRS7, TRS12, TRS14, TRS20, TRS25, TRS40, and TRS43, in order to select RAPD primers useful for

analysis. Amplicons were obtained for 482 primers, with an average of eight amplicons of different sizes identified for each primer. Based on electrophoretic evaluation of amplification pattern quality, 68 primers that produced polymorphic amplicons were chosen for the assessment of genetic diversity of the *T. atroviride* strains. RAPD PCR was performed with selected primers on 38 strains of *T. atroviride* collected at different locations in Poland (Table 1). As a control, *T. atroviride* reference strains CBS 693.94 (Dodd et al. 2003) and IMI 206040 (Kubicek et al. 2011) were used. High-quality amplification products were obtained for 55 of the 68 RAPD primers. Only two primers (OPK11 and OPX09) did not show any polymorphisms among tested strains. The greatest number of polymorphic amplicons distinguishing groups of strains was obtained with primer OPA10. Amplification profiles of three primers, OPI18, OPQ01, and OPZ04, showed that the *T. atroviride* strains, with the exception of strain TRS18, can be classified into two main groups. The amplification profile generated using OPZ04 is shown in Fig. 2.

RAPD amplicons were scored for presence and absence, and 391 amplicons differentiating tested strains were used to construct a binary matrix, and then a dendrogram illustrating genetic diversity (Fig. 3). The cophenetic correlation coefficient ( $r$ ) for the dendrogram was 0.979, indicating a good fit between the unmolded data and the dendrogram. Tested *T. atroviride* strains, of which all except TRS18 belonged to clade A, were classified into two main groups. The first group was represented by the reference strain IMI206040, as well as 28 of the tested strains, and corresponds to the two biggest groups of strains including TRS43, distinguished based on *tef1* sequence. The second group contained reference strain CBS 693.94 and the following eight tested strains: TRS2, TRS6, TRS10, TRS16, TRS26, TRS40, TRS42, and TRS44. This group corresponds to CBS 693.94 and remaining two small groups of strains distinguished based on *tef1*. The observed variation among strains within the two major groups was rather limited; however, some subgroups could be distinguished (Fig. 3). *T. atroviride* strain TRS18 was clearly distinct from both groups of strains, based on both *tef1* and RAPD analysis, and also did not fit any previously described clades. TRS18 *tef1* intron 4 sequence BLAST similarity search with non-redundant nucleotide sequence database at NCBI revealed 95 % identity with any sequence deposited in GenBank. This suggests that TRS18 strain is characterized by a unique and distinct genotype that could represent new clade of *T. atroviride*.

Twenty polymorphic RAPD markers were chosen for cloning, and 13 RAPD-PCR products were successfully cloned and sequenced (Table 2). Resulting sequences were aligned with the *T. atroviride* IMI206040 genome at the JGI portal. Twelve RAPD sequences were identical to corresponding *T. atroviride* IMI206040 sequences (e-value=0). Sequence




**Fig. 1** Bayesian identification phylogram of *T. atroviride* strains used in this study based on *tef1* gene sequence alignments. Numbers at nodes indicate the posterior probability coefficients obtained after one million generations. IMI206040 and CBS693.94 are reference strains of *T. atroviride*


R08-1283 showed some variation, but the e-value was very low, indicating strong similarity to the *T. atroviride* genome (Table 2). Seven sequences showed strong similarity to hypothetical proteins of *T. atroviride* deposited in the NCBI database, and Pfam protein domains were clearly identified for three of these (Table 2). The H08-917 sequence showed a high degree of similarity to tetratricopeptide repeat (TPR)-containing proteins. TPR motifs are short amino acid repeats that occur in many proteins and not only are responsible for interactions between proteins but also facilitate protein complex formation and protein transport within cells. TPR-containing proteins play a role in protein assembly and therefore have a significant impact on the cell cycle and a plethora of cellular processes (D'Andrea and Regan 2003). Interestingly, the H08-917 sequence also possessed short and degenerate DNA repeats that were identified by RepeatMasker and mfold. The R08-1283 and T07-429

sequences were recognized as proteins with ankyrin (ANK) repeats, which play a role in protein-protein interactions (Rubtsov and Lopina 2000). Moreover, the X16-1030 sequence was similar to the NIPA membrane protein, which functions as a magnesium transporter (Goytain et al. 2007, 2008).

Using the identified RAPD marker sequences and the *T. atroviride* genome sequence, pairs of specific primers were designed, followed by optimization of PCR conditions and testing of the primers on a set of *Trichoderma* strains belonging to different species, including *T. atroviride* (Table 3). Unique PCR amplification products for three pairs of primers, Q01-4, X18-35, and Z04-2, were obtained for *T. atroviride* strains and not for 16 other *Trichoderma* species/clades (Fig. 4). In the case of X18-35, a very faint PCR amplicon was obtained for *T. gamsii*. These suggest that the amplicons Q01-4, X18-35, and Z04-2 are specific to *T. atroviride*. We



**Fig. 2** RAPD-PCR amplification profile generated with primer OPZ04 for *T. atroviride* strains used in this study. Polymorphic amplicon OPZ04-873 bp is shown by the arrow. IMI206040 and CBS 693.94 are reference strains of *T. atroviride*. M, 1-kb DNA size marker




**Fig. 3** Dendrogram generated using Jaccard's coefficient and the UPGMA clustering method. Binary matrices for *T. atroviride* strains were constructed based on evaluation of RAPD amplicons generated using 55 primers. Cluster analysis of the binary data was performed

using NTSYS-pc 2.1 software. Similarity matrices were generated using Jaccard's coefficient and an unweighted pair-group method using arithmetic averages (UPGMA) was used to generate the dendrogram

propose that pairs of primers Q01-4, Z18-35, and Z04-2 could be used for *T. atroviride* detection.


For selected *T. atroviride* strains representing two main groups of strains, amplicons Q01-4, X18-35, and Z04-2 were

sequenced, and sequences were deposited in GenBank (Table 3). These sequences were compared, and polymorphic restriction sites were identified. Based on identified sequence differences (single nucleotide polymorphisms and insertions/



**Fig. 4** Amplification of Q01-4, X18-35, and Z04-2 fragments in different *Trichoderma* species. M, DNA size markers; amplicon 5.8S-R-LR6, PCR positive control





**Fig. 5** Q01-4 and Z04-2 CAPS marker profiles generated by PCR amplification with primers Q01\_4F and Q01\_4R and digestion of PCR products with *Bsl*I (upper part) and primers Z04\_2F and Z04\_2R and

digestion of PCR products with *Dra*I. IMI206040 and CBS 693.94 are the reference strains of *T. atroviride*. CAPS, cleaved amplified polymorphic sequences. M, 100-bp Plus DNA ladder

deletions), restriction enzymes were selected to digest PCR amplicons and develop CAPS markers useful for *T. atroviride* identification (Table 3). In the case of CAPS marker Q01-4 (Q01-4FR amplification product digested with *Bsl*I), a two-band pattern was obtained for 29 strains of *T. atroviride* belonging to group 1, and a single band corresponding to the non-restricted Q01-4 F-4R amplicon was obtained for group 2 strains and TRS18 (Fig. 5). This CAPS marker allows the distinction of strains belonging to *T. atroviride* group 1. For Z04-2 amplification products, *Dra*I digestion resulted in banding patterns reflecting the RAPD-based grouping. However, the products for strain TRS6 and TRS18 were digested and these strains were not classified to the group 1 as in the RAPD study (Fig. 5). We propose that analysis of both CAPS markers can be used to extend ITS/*tef1*-based *T. atroviride* identification and can assist in classification of strains within clade A into one of the two main groups of *T. atroviride*: group 1 (IMI206040-like), characterized by both Q01-4 and Z04-2 amplicons digested, and group 2 (CBS 693.60-like), characterized by both amplicons non-digested. In the case where Q01-4 amplicon is not digested and Z04-2 is digested, additional markers have to be developed to distinguish TRS6-like from TRS18-like strains. Developed CAPS markers have to be further tested on a larger number of strains that will confirm their usefulness in *T. atroviride* genetic studies. The variable RAPD loci found in this study and characterized at the sequence level may be useful in the development of more complex assays, such as

multiplex PCR, real-time PCR, or next-generation sequencing (NGS)-based assays, that will be valuable in identification and monitoring of *T. atroviride* strains, as well as in genetic diversity studies including metagenomics approaches and further *T. atroviride* pan-genome understanding.

**Acknowledgments** This study was conducted as a part of the project “Polish *Trichoderma* strains in plant protection and organic waste management” under Priority 1.3.1, subject area “Bio,” co-financed by The European Union through the European Regional Development Fund within the Innovative Economy Operational Program, 2007–2013, project No UDA-POIG.01.03.01-00-129/09-08. The authors declare that there is no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

- Abbasi PA, Miller SA, Meulia T, Hoitink HAJ, Kim J-M (1999) Precise detection and tracing of *Trichoderma hamatum* 382 in compost-amended potting mixes by using molecular markers. *Appl Environ Microbiol* 65:5421–5426
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402

- Atanasova L, Druzhinina IS (2010) Global nutrient profiling by Phenotype MicroArrays: a tool complementing genomic and proteomic studies in conidial fungi. *J Zhejiang Univ Sci B (Biomed and Biotechnol)* 11:151–168
- Atanasova L, Druzhinina IS, Jaklitsch WM (2013) Two hundred *Trichoderma* species recognized on the basis of molecular phylogeny. In: Mukherjee PK, Horwitz BA, Singh US, Mukherjee M, Schmoll M (eds) *Trichoderma: biology and applications*. CABI, Wallingford, pp 10–42
- Benítez T, Rincón AM, Limón MC, Codón AC (2004) Biocontrol mechanisms of *Trichoderma* strains. *Int Microbiol* 7:249–260
- Błaszczak L, Popiel D, Chełkowski J, Koczyk G, Samuels GJ, Sobieralski K, Siwulski M (2011) Species diversity of *Trichoderma* in Poland. *J Appl Genet* 52:233–243
- Bowen K, Francevic SC, Crowhurst RN, Templeton MD, Stewart A (1996) Differentiation of a specific *Trichoderma* biological control agent by restriction fragment length polymorphism (RFLP) analysis. *N Z J Crop Hortic* 24:207–217
- Buhariwalla HK, Sriakashmi P, Kannan S, Kanchi RS, Chandra S, Satyaprasad K, Waliyar F, Thakur RP, Crouch JH (2005) AFLP analysis of *Trichoderma* spp. from India compared with sequence and morphological-based diagnostics. *J Phytopathol* 153:389–400
- Carbone I, Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous *Ascomycetes*. *Mycologia* 91:553–556
- Chakraborty BN, Chakraborty U, Sunar K, Dey PL (2011) RAPD profile and rDNA sequence analysis of *Talaromyces flavus* and *Trichoderma* species. *Indian J Biotechnol* 10:487–495
- Chen X, Romaine CP, Tan Q, Schlaghauser B, Ospina-Giraldo MD, Royse DJ, Huff DR (1999) PCR-based genotyping of epidemic and preepidemic *Trichoderma* isolates associated with green mold of *Agaricus bisporus*. *Appl Environ Microbiol* 65:2674–2678
- Chet I, Inbar J (1994) Biological control of fungal pathogens. *Appl Biochem Biotechnol* 48:37–43
- Cooke DEL, Duncan JM (1997) Phylogenetic analysis of *Phytophthora* species based on ITS1 and ITS2 sequences of the ribosomal RNA gene repeat. *Mycol Res* 101:667–677
- Cordier C, Edel-Hermann V, Martin-Laurent F, Blal B, Steinberg C, Alabouvette C (2007) SCAR-based real time PCR to identify a biocontrol strain (T1) of *Trichoderma atroviride* and study its population dynamics in soils. *J Microbiol Methods* 68:60–68
- D'Andrea LD, Regan L (2003) TPR proteins: the versatile helix. *Trends Biochem Sci* 28:655–662
- Devi TP, Kamil D, Prabhakaran N (2011) Development of genus specific rDNA based marker for detection of *Trichoderma* species. *J Mycol Plant Pathol* 41:600–604
- Dodd SL, Lieckfeldt E, Samuels GJ (2003) *Hypocrea atroviridis* sp. nov., the teleomorph of *Trichoderma atroviride*. *Mycologia* 95:27–40
- Dodd SL, Hill RA, Stewart A (2004a) Monitoring the survival and spread of the biocontrol fungus *Trichoderma atroviride* (C65) on kiwifruit using a molecular marker. *Australas Plant Pathol* 33:189–196
- Dodd SL, Hill RA, Stewart A (2004b) A duplex-PCR bioassay to detect a *Trichoderma virens* biocontrol isolate in non-sterile soil. *Soil Biol Biochem* 36:1955–1965
- Druzhinina I, Kubicek CP (2005) Species concepts and biodiversity in *Trichoderma* and *Hypocrea*: from aggregate species to species clusters? *J Zhejiang Univ Sci B* 6:100–112
- Druzhinina IS, Kopchinskiy AG, Komoń M, Bissett J, Szakacs G, Kubicek CP (2005) An oligonucleotide barcode for species identification in *Trichoderma* and *Hypocrea*. *Fungal Genet Biol* 42:813–828
- Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV, Kubicek CP (2011) *Trichoderma*: the genomics of opportunistic success. *Nat Rev Microbiol* 9:749–759
- Feng XM, Holmberg A-IJ, Sundh I, Ricard T, Melin P (2011) Specific SCAR markers and multiplex real-time PCR for quantification of two *Trichoderma* biocontrol strains in environmental samples. *BioControl* 56:903–913
- Friedl MA, Druzhinina IS (2012) Taxon-specific metagenomics of *Trichoderma* reveals a narrow community of opportunistic species that regulate each other's development. *Microbiology* 158:69–83
- Gal-Hemed I, Atanasova L, Komon-Zelazowska M, Druzhinina IS, Viterbo A, Yarden O (2011) Marine isolates of *Trichoderma* spp. as potential halotolerant agents of biological control for arid-zone agriculture. *Appl Environ Microbiol* 77:5100–5109
- Goytain A, Hines RM, El-Husseini A, Quamme GA (2007) NIPA1 (SPG6), the basis for autosomal dominant form of hereditary spastic paraplegia, encodes a functional  $Mg^{2+}$  transporter. *J Biol Chem* 282:8060–8068
- Goytain A, Hines RM, Quamme GA (2008) Functional characterization of NIPA2, a selective  $Mg^{2+}$  transporter. *Am J Physiol Cell Physiol* 295:C944–C953
- Grigoriev IV, Nordberg H, Shabalov I, Aerts A, Cantor M, Goodstein D, Kuo A, Minovitsky S, Nikitin R, Ohm RA, Otilar R, Poliakov A, Ratnere I, Riley R, Smirnova T, Rokhsar D, Dubchak I (2012) The genome portal of the department of energy joint genome institute. *Nucleic Acids Res* 40:D26–D32
- Hahn A, Wallisch S, Radl V, Charles Munch J, Schlöter M (2007) A new cultivation independent approach to detect and monitor common *Trichoderma* species in soils. *J Microbiol Methods* 69:86–92
- Harman GE, Howell CR, Viterbo A, Chet I, Lorito M (2004) *Trichoderma* species — opportunistic, avirulent plant symbionts. *Nat Rev Microbiol* 2:43–56
- Hermosa MR, Grondona I, Iturriaga EA, Diaz-Minguez JM, Castro C, Monte E, Garcia-Acha I (2000) Molecular characterization and identification of biocontrol isolates of *Trichoderma* spp. *Appl Environ Microbiol* 66:1890–1898
- Hermosa MR, Grondona I, Diaz-Minguez JM, Iturriaga EA, Monte E (2001) Development of a strain-specific SCAR marker for the detection of *Trichoderma atroviride* 11, a biological control agent against soilborne fungal plant pathogens. *Curr Genet* 38:343–350
- Hoyos-Carvajal L, Orduz S, Bissett J (2009) Genetic and metabolic biodiversity of *Trichoderma* from Colombia and adjacent neotropical regions. *Fungal Genet Biol* 46:615–631
- Jaklitsch WM, Komon M, Kubicek CP, Druzhinina IS (2005) *Hypocrea voglmayrii* sp. nov. from the Austrian Alps represents a new phylogenetic clade in *Hypocrea/Trichoderma*. *Mycologia* 97:1365–1378
- Jaklitsch WM, Samuels GJ, Dodd SL, Lu B-S, Druzhinina IS (2006) *Hypocrea rufa/Trichoderma viride*: a reassessment, and description of five closely related species with and without warted conidia. *Stud Mycol* 56:135–177
- Kindermann J, El-Ayouti Y, Samuels GJ, Kubicek CP (1998) Phylogeny of the genus *Trichoderma* based on sequence analysis of the internal transcribed spacer region 1 of the rDNA cluster. *Fungal Genet Biol* 24:298–309
- Kopchinskiy A, Komoń M, Kubicek CP, Druzhinina IS (2005) *TrichoBLAST*: a multilocus database for *Trichoderma* and *Hypocrea* identifications. *Mycol Res* 109:658–660
- Kredics L, Kocsubé S, Nagy L, Komoń-Zelazowska M, Manczinger L, Sajben E, Nagy A, Vágvolgyi C, Kubicek CP, Druzhinina IS, Hatvani L (2009) Molecular identification of *Trichoderma* species associated with *Pleurotus ostreatus* and natural substrates of the oyster mushroom. *FEMS Microbiol Lett* 300:58–67
- Kredics L, Hatvani L, Naeimi S, Körmöczy P, Manczinger L, Vágvolgyi C, Druzhinina I (2014) Biodiversity of the genus *Hypocrea/Trichoderma* in different habitats. In: Gupta VK, Schmoll M, Herrera-Estrella A, Upadhyay RS, Druzhinina I, Tuohy MG (eds) *Biotechnology and biology of Trichoderma*. Elsevier, London, pp 3–24

- Kubicek CP, Bissett J, Druzhinina IS, Kullnig-Gradinger C, Szakacs G (2003) Genetic and metabolic diversity of *Trichoderma*: a case study on South-East Asian isolates. *Fungal Genet Biol* 38:310–319
- Kubicek CP, Herrera-Estrella A, Seidl-Seiboth V, Martinez DA, Druzhinina IS, Thon M, Zeilinger S, Casas-Flores S, Horwitz BA, Mukherjee PK, Mukherjee M, Kredics L, Alcaraz LD, Aerts A, Antal Z, Atanasova L, Cervantes-Badillo MG, Challacombe J, Chertkov O, McCluskey K, Couplier F, Deshpande N, von Döhren H, Ebbole DJ, Esquivel-Naranjo EU, Fekete E, Flippin M, Glaser F, Gómez-Rodríguez EY, Gruber S, Han C, Henrissat B, Hermosa R, Hernández-Oñate M, Karaffa L, Kosti I, Le Crom S, Lindquist E, Lucas S, Lübeck M, Lübeck PS, Margeot A, Metz B, Misra M, Nevalainen H, Omann M, Packer N, Perrone G, Uresti-Rivera EE, Salamov A, Schmoll M, Seiboth B, Shapiro H, Sukno S, Tamayo-Ramos JA, Tisch D, Wiest A, Wilkinson HH, Zhang M, Coutinho PM, Kenerley CM, Monte E, Baker SE, Grigoriev IV (2011) Comparative genome sequence analysis underscores mycoparasitism as the ancestral life style of *Trichoderma*. *Genome Biol* 12:R40
- Kullnig CM, Krupica T, Woo SL, Mach RL, Rey M, Benítez T, Lorito M, Kubicek CP (2001) Confusion abounds over identities of *Trichoderma* biocontrol isolates. *Mycol Res* 105:769–772
- Longa CMO, Savazzini F, Tosi S, Elad Y, Pertot I (2009) Evaluating the survival and environmental fate of the biocontrol agent *Trichoderma atroviride* SC1 in vineyards in northern Italy. *J Appl Microbiol* 106:1549–1557
- Lorito M, Woo SL, Harman GE, Monte E (2010) Translational research on *Trichoderma*: from ‘omics to the field. *Annu Rev Phytopathol* 48:1–23
- Miyazaki K, Tsuchiya Y, Okuda T (2009) Specific PCR assays for the detection of *Trichoderma harzianum* causing green mold disease during mushroom cultivation. *Mycoscience* 50:94–99
- Mulaw TB, Kubicek CP, Druzhinina IS (2010) The rhizosphere of *Coffea arabica* in its native highland forests of Ethiopia provides a niche for a distinguished diversity of *Trichoderma*. *Diversity* 2:527–549
- Naef A, Senatore M, Défago G (2006) A microsatellite based method for quantification of fungi in decomposing plant material elucidates the role of *Fusarium graminearum* DON production in the saprophytic competition with *Trichoderma atroviride* in maize tissue microcosms. *FEMS Microbiol Ecol* 55:211–220
- Naeimi S, Kocsubé S, Antal Z, Okhovvat SM, Javan-Nikkhah M, Vágvolgyi C, Kredics L (2011) Strain-specific SCAR markers for the detection of *Trichoderma harzianum* AS12-2, a biological control agent against *Rhizoctonia solani*, the causal agent of rice sheath blight. *Acta Biol Hung* 62:73–84
- Olczak-Woltman H, Bartoszewski G, Madry W, Niemirówicz-Szczytt K (2009) Inheritance of resistance to angular leaf spot (*Pseudomonas syringae* pv. *lachrymans*) in cucumber and identification of molecular markers linked to resistance. *Plant Pathol* 58:145–151
- Punta M, Coghill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer ELL, Eddy SR, Bateman A, Finn RD (2012) The Pfam protein families database. *Nucleic Acids Res* 40:D290–D301
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* 61:539–542
- Rubio MB, Hermosa MR, Keck E, Monte E (2005) Specific PCR assays for the detection and quantification of DNA from the biocontrol strain *Trichoderma harzianum* 2413 in soil. *Microb Ecol* 49:25–33
- Rubtsov AM, Lopina OD (2000) Ankyrins. *FEBS Lett* 482:1–5
- Samuels GJ, Dodd SL, Lu BS, Petrini O, Schroers HJ, Druzhinina IS (2006) The *Trichoderma koningii* aggregate species. *Stud Mycol* 56:67–133
- Savazzini F, Longa CMO, Pertot I, Gessler C (2008) Real-time PCR for detection and quantification of the biocontrol agent *Trichoderma atroviride* strain SC1 in soil. *J Microbiol Methods* 73:185–194
- Schuster A, Schmoll M (2010) Biology and biotechnology of *Trichoderma*. *Appl Microbiol Biotechnol* 87:787–799
- Stewart A, Hill R (2014) Applications of *Trichoderma* in plant growth promotion. In: Gupta VK, Schmoll M, Herrera-Estrella A, Upadhyay RS, Druzhinina I, Tuohy MG (eds) *Biotechnology and biology of Trichoderma*. Elsevier, London, pp 415–428
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J Bacteriol* 172:4238–4246
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, pp 315–322
- Woo SL, Scala F, Ruocco M, Lorito M (2006) The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology* 96:181–185
- Zimand G, Valinsky L, Elad Y, Chet I, Manulis S (1994) Use of the RAPD procedure for the identification of *Trichoderma* strains. *Mycol Res* 98:531–534
- Zuker M (2003) Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* 31:3406–3415